

## Molecular Analysis on *Streptococcus Mutans*, In Detecting the Mutacin I and II and GTF Genes in Caries Free and Caries Active Individuals

Sanjay Rathod<sup>1</sup>, C.T. Shivannaveur<sup>2</sup>, Raghvendre Ramachari<sup>3</sup>, Suresh Kumar M.<sup>4</sup>

### Abstract

*Streptococcus mutans* is a major pathogen implicated in dental caries. Its virulence is enhanced by its ability to produce bacteriocins, called mutacins, which inhibit the growth of other Gram-positive bacteria. Sucrose-dependent adherence is mediated by glucans, polymers of glucose synthesized from sucrose by glucosyltransferase (Gtf) enzymes. *S. mutans* makes several proteins that have the property of binding glucans. We hypothesized that three of these glucan-binding proteins (Gbps), Gbps A, C and D, contribute to the carcinogenicity of *S. mutans*. The availability of suitable in vitro and in vivo systems and a thorough understanding of the genetics, biochemistry and physiology of the dental pathogen *Streptococcus mutans* have greatly advanced in understanding of important areas in the field of bacteriology such as interspecies biofilms, competence development and stress responses. *S. mutans*, as an organism had evolved in close association with the human host, as a novel Gram-positive model organism. The role of mutacins *in vivo* is unclear, however the antimicrobial activity of these substances may confer an ecological advantage for the producing strain in bacterial communities such as dental biofilm (56), and they may also be important for the establishment of *S. mutans in vivo*. Mutacins are peptide or protein antibiotics that are mainly bactericidal for other bacteria of the same or closely related species, as well as for other Gram-positive microorganisms, and are likely to confer an ecological advantage in diverse bacterial communities such as dental biofilm. The relationship between caries activity and the higher synthesis of some virulence factors by different genotypes of *S. mutans* has been related. In this study, of 17 isolates, 10 isolates from caries-active and 7 isolates from caries-free were randomly selected for the molecular analysis for the mutacin I and II and for the (GTF) genes studies. It is concluded in the study that Mutacin I and II and (GTF) enzymes detected contribute to the carcinogenicity of *S. mutans* through a mechanism that may involve alteration of biofilm architecture. In addition, reports of higher numbers of *S. mutans* genotypes with increased virulence in caries-active Subjects suggest the importance of microenvironmental factors in increasing the risk of caries active individuals.

**Keywords:** S. Mutans; Caries; Biofilm; Glucan; Mutacin I and II; Glycosyltransferase Gene (GTF).

### Introduction

Dental caries is a transmissible infectious disease in which mutans streptococci (MS) play the major role. As in many infectious diseases, colonization by pathogens is required before the disease can occur. MS are generally considered to be the principal etiological agent of dental Caries. There is a range of virulence factors important for the establishment of

MS in the complex microbial community of dental biofilm. Studies of the virulence factors of *S. mutans* and their correlation with species biodiversity are fundamental to understanding the role played by colonization by different genotypes in the same individual, and the expression of characteristics that may or may not influence their virulence capacity and survival ability under different environmental conditions. In 1924, J. Clarke isolated an organism

<sup>1</sup>Guest Lecturer, <sup>2</sup>Professor, <sup>3,4</sup>Research Scholar, Department of Microbiology, Gulbarga University, Gulbarga -585106, Karnataka.

from carious lesions and called it *S. mutans*, because he thought the oval-shaped cells observed were mutant forms of streptococci (Clarke, 1924). However, it was only in the late 1950s that *S. mutans* received greater attention from the scientific community and, by the mid 1960s, it was recognized as a major aetiological agent in dental caries (Loesche, 1986). In the subsequent two decades, researchers began to uncover the pathophysiology of *S. mutans*.

The major virulence traits of *S. mutans* were established: (i) the ability to produce large quantities of organic acids (acidogenicity) from metabolized carbohydrates; (ii) the ability to survive at low pH (aciduricity); and (iii) the ability to synthesize extracellular glucan-homopolymers from sucrose, which play a critical role in initial attachment, colonization and accumulation of biofilms on tooth surfaces (Banas & Vickerman, 2003; Bowen & Koo, 2011; Burne, 1998; Loesche, 1986). With the advances in molecular genetics techniques in the 1980s and 90s, scientists began to more rapidly understand how metabolic pathways enabled *S. mutans* to evolve into a specialized dental pathogen.

In this study we have, selected *S. mutans* isolates from both with caries active and caries free individuals, to know the types of mutacin genes Present. Using specific primers for *mut I* and *mut II* genes and genomic DNA from *S. mutans* isolates amplified 750 bp and 444bp amplicons by simple PCR method. Amplicons for *mut I* and *mut II* genes generated by each isolate. All *S. mutans* isolates from caries individuals express the *mut I* (750 bp) compared to only 70% of isolates from caries free individuals tested for molecular study. However, *mut II* (444 bp) gene is not amplified in all the isolates tested. the glucosyltransferase (GTF) enzymes is considered fundamental for the virulence of *S. mutans* in the pathogenesis of dental caries. *S. mutans* in the presence of sucrose produces this enzyme which converts sucrose to water insoluble glucan (WIG) which is directly proportional to the plaque formation *in vivo* and biofilm *in vitro*. Significant association between the presence of *gtfB*, production of adhesive substances and percentages of growing bacteria adhering to glass surfaces in the presence of sucrose has been observed.

The percent adherence of *S. mutans* isolates from with caries individuals is higher (39.56%) than the *S. mutans* isolated from without caries (19.11%) individuals These results clearly indicate that *S. mutans* isolated from with caries individuals are having more capacity to form dental plaques than from caries free individuals.

## Materials and Methods

The study was planned to investigate the prevalence of the bacterium present in the oral cavity (saliva) which plays a principle role in the initiation of dental caries. Factors affecting the prevalence of bacterium and its virulence in caries risk individuals. We selected two dental college hospitals in Gulbarga namely, Nijalingappa Dental College and Hospital and Al-Bader's Dental College and Hospital, for the collection of saliva samples from subjects for this study.

The study comprised of a total 254 subjects, which included both the caries active and caries free individuals with or without tobacco chewing habit volunteers attending the outpatient department (OPD) of two dental colleges and hospitals of Gulbarga city. They were selected in such a way that they belonged to different sex, age and socio-economic background. Unstimulated 2–3 ml of saliva sample was collected into a pre-labeled sterile wide mouthed plastic capped bottles from the volunteer individuals with or without caries and the collected saliva samples were immediately transported to the laboratory and processed for the enumeration of mutant streptococci and isolation of *S. mutans*. The saliva collected from each individual processed for the total count of mutans streptococci and *S. mutans* by serially diluting 1 ml of well-mixed saliva up to  $10^{-5}$  dilutions in sterile normal saline. From the  $10^{-2}$  to  $10^{-6}$  dilutions, 0.5 ml from each tube were inoculated on to Mitis Salivarius Bacitracin (MSB) agar media by spread plate method and plates were incubated in an anaerobic jar for 48 hrs. The plates were observed for the growth and the colonies were counted and the bacterial load in the collected saliva sample was calculated.

### Composition of Mitis-Salivarius Bacitracin (MSB) Agar Medium

Component	gms/litre.
Casein enzymatic	15.00
Hydro lysate peptic	5.00
Digest of animal tissue	1.00
Dextrose	50.00
Sucrose	50.00
Dipotassium phosphate	4.00
Tryphan blue	0.075
Crystal violet	0.0008
Water	1000.0 ml
Agar	15.00
Final pH = 7.0 ± 0.2	

The colony showing distinct cultural morphological characteristics of *S. mutans* was subcultured and growth was used for the performance of biochemical tests. Rapid biochemical identification kit from HI-Media was used along with conventional biochemical tests like fermentation of mannitol and sorbitol, hydrolysis of arginine and esculin and Voges Proskauer. Based on the cultural, biochemical and morphological characteristics the isolates were identified up to the species level. The rapid biochemical identification kit was purchased from the Hi-media Pvt. Ltd., Mumbai and other biochemical tests were prepared and Performed according to the procedures described in the standard microbiology laboratory manuals.

#### *Molecular Studies of S. mutans Isolates*

As dental caries is a microbial disease so the prerequisite for caries development is the presence of dental plaque on the teeth and unless this biofilm present caries will not occur, regardless of any other risk factors (Kidd, 1999). So, initiation and progress of the dental caries mainly based on the survivability of *S. mutans* in saliva and especially in the dental plaque (biofilm) by producing adherence substances like glucan and bacteriocins like mutacins. In order to understand and know the cariogenicity of the isolated *S. mutans*, we selected total of 17 *S. mutans* isolates 10 from without caries and 7 from with caries individuals for molecular typing of *gtf-B* and mutacin genes by PCR amplification.

#### *PCR Amplification of gtf B Gene*

In this study, the 517 bp region off *gtf B* was amplified using the following primers by simple PCR (Oho et al., 2000):

*gtfB* Forward 5'-ACTACACTTTCGGGTGGCTTGG-3'  
*gtfB* Reverse 5'-CAGTATAAGCGCCAGTTCATC-3'

*Conditions for Amplification* (Oho et al., 2000): For the PCR running standardized the condition used by the Oho et al., 2000; in the Clinspire Technology, Bellary, Karnataka. The conditions applied for the amplification of *gtf B* gene using eppendorf gradient PCR machine are as follows:

#### *Conditions for Gtf B Amplification*

Initial denaturation step: at 95°C for 5 min.  
 For each cycle (total 30 cycles)  
 Followed by denaturation: 95°C for 30 seconds,

Annealing at : 59°C for 30 seconds  
 Extension at : 72°C for 1 min  
 Final elongation at : 72°C for 10 min and stand at 10°C

#### *PCR Amplification of Mutacin Genes*

*Primers:* We used the following primers for amplification of Type-I (750bp) and Type- II (444bp) by PCR.

*Mutacin: Type-I* (Qi et al., 1999b)

Forward 5'-AGTTTCAATAGTTACTGTTGC-3'  
 Reverse 5'-GCCAAACGGAGTTGATCTCGT-3'

*Mutacin - Type-II* (Novak et al., 1994)

Forward 5'- AACGCAGTAGTTTCTTTGAA-3'  
 Reverse 5'- TTCCGGTAAGTACATAGTGC-3'

*PCR Conditions for Mutacin Gene Amplification* (Qi et al., 2001; Kamiay et al., 2005):

Initial denaturation step : at 94°C for 5 min.

For each cycle (total 35 cycles)

Followed by denaturation : 94°C for 45 seconds,

Annealing at : 52°C for 1 min

Extension at : 72°C for 2 min

Final elongation at : 72°C for 7 min and stand at 10°C.

## **Results and Discussion**

A total of 120 (47.24%) *S. mutans* was isolated from 254 saliva samples collected without stimulant from with caries and caries free individuals with and without habitual Incidence rate was very high (57.04%) among the individuals with caries as compared to individuals without caries (34.82%). Incidence of *S. mutans* in non habituals was observed to be 55.03% (82/149) which is much higher than that in the individuals with habits (36.19%; 38/105), however it is maximum (63.46%) in the habitual males with caries compared to any group in this study. We observed higher (60.71%) incidence of *S. mutans* in habitual individuals with caries and lowest (08.16%) in habitual individuals without caries. Over all higher incidents observed in female (51.54%) than in male (44.58%), however it is reverse among genders with caries individuals.

### Molecular Studies of *S. mutans* Isolates

Mutacin I, the focus of this study, is a 24 aa lantibiotic with a molecular mass of 2364 Da. The mutacin I biosynthesis gene locus consists of 14 genes in the order of *mutR*, *mutA*, *mutA9*, *mutB*, *mutC*, *mutD*, *mutP*, *mutT*, *mutF*, *mutE*, *mutG*, *orfX*, *orfY*, *orfZ* (Qi et al., 2000). *MutR* is thought to be the positive regulator for the expression of the mutacin I operon. *MutA* and *MutA9* show strong similarity to each other. While *mutA* is the structural gene for prepromutacin I, *mutA9* is not required for mutacin I activity. *MutB*, *MutC* and *MutD* constitute the modification apparatus for the premature peptide, and *MutT* and *MutP* are the ABC transporter and protease, respectively, for the transporting and processing of premature mutacin. *MutF*, *MutE* and *MutG* are probable immunity proteins for mutacin I. The functions of *OrfX*, *OrfY* and *OrfZ* proteins in mutacin I production are unknown (Qi et al., 2000) (*gtf*) enzymes contribute to the ability of *S. mutans* to induce caries in the presence of sucrose. Of the individual Gbp mutants, only the *gbpD* strain was attenuated relative to the parental. We propose that the combined loss of Gbps A and C has the most dominant effect on *S. mutans* cariogenicity, though the additive loss of GbpD extends the magnitude and breadth of the attenuation. While the loss of GbpD alone mostly affects caries development on sulcal surfaces, the combined loss of Gbps A and C has its greatest impact on smooth (buccal) Surfaces. Clearly, each Gbp makes a unique contribution to the caries process, but the mechanistic contributions of each cannot be explained by the *in vivo* results alone. We can only speculate why the *gbpD* mutant is attenuated but the *gbpAD* and *gbpCD* mutants are not. It would appear that the loss of either GbpA or GbpC in some way compensates for the loss of GbpD, but when paired, the *gbpAC* mutant is significantly less cariogenic. Evidence supporting regulated expression of GbpC was provided by Biswas et al. [33], where expression of *gbpC* mRNA was observed to peak at mid-log phase and was extremely diminished in stationary phase cultures.

As dental caries is a microbial disease so the prerequisite for caries development is the presence of dental plaque on the teeth and unless this biofilm present caries will not occur, regardless of any other risk factors (Kidd, 1999). So, initiation and progress of the dental caries mainly based on the survivability of *S. mutans* in saliva and especially in the dental plaque (biofilm) by producing adherence substances like glucan and bacteriocins like mutacins. In order to understand and know the cariogenicity of the isolated *S. mutans*, we selected total of 17 *S. mutans*

isolates 10 from without caries and 7 from with caries individuals for molecular typing of *gtf-B* and mutacin genes by PCR amplification.




**Fig. 1:** *Streptococcus mutans* growth on the mitis salivarius bacitracin (MSB) agar

**Table 1:** Over all incidence of *Streptococcus mutans* in saliva samples

		No. of Samples			Incidence of <i>S.Mutans</i>		
		male	female	Total	male	female	Total
With Caries	Habitual	52	04	56	33 (63.46%)	01 (25.00%)	34 (60.71%)
	Non-habitual	38	48	86	20 (52.63%)	27 (56.25%)	47 (54.65%)
Total		90	52	142	53 (58.88%)	28 (53.84%)	81 (57.04%)
Without Caries	Habitual	43	06	49	03 (06.97%)	01 (16.66%)	04 (08.16%)
	Non-Habitual	24	39	63	14 (54.16%)	21 (53.84%)	35 (34.82%)
Total		67	45	112	17 (25.37%)	22 (48.88%)	39 (34.24%)
Grand Total		157	97	254	70 (44.58%)	50 (51.54%)	120 (47.24%)



**Fig. 2:** Mutacin genes mut I and II in *S. mutans*.  
Lane M-250 bp DNA ladder  
Lane 1 to 10: mut I gene amplicons of *S. mutans* isolates from without caries individuals  
Lane 11 to 17: mut I gene amplicons of *S. mutans* isolates from with caries individuals  
Lane 18 to 24: mut II gene amplicons of *S. mutans* isolates from with caries individuals shows no amplicons



**Fig. 3:** GTF B gene amplicons of *S. mutans* isolates.  
Lane M-100 bp DNA ladder  
Lane 1 to 7: *S. mutans* isolates from with caries and  
Lane 8 to 10: *S. mutans* isolates from without caries individuals

### Conclusion

1. Amplicons of *gtf B* gene from all *S. mutans* isolates tested showed single band on agarose gel corresponding to 517 bp region. Study suggested that isolated streptococci are *S. mutans*

and cariogenic in nature.

2. Higher adherence capacity of *S. mutans* isolates from with caries individuals to glass surface and presence of *mut I* gene in all *S. mutans* isolates from caries individuals suggesting the role of *gtf B* and *mut I* in the cariogenicity of *S. mutans*.
3. From the molecular studies of *S. mutans* isolates from saliva collected from caries individuals exhibited higher cariogenic virulence factors than *S. mutans* from caries free individuals. Our study emphasizes the presence of virulence factors in the cariogenic strains of *S. mutans* as essential to initiate the dental caries in the susceptible individuals of any age group.
4. Further research on the formation of biofilm in animal models and in *in vitro* by the *S. mutans* isolated from different niche of oral cavity of caries and caries free individuals of different geographical areas is required to confirm the exact role of *gtf* and *mut* genes in the cariogenesis of *S. mutans*.
5. Molecular study of virulent factors of *S. mutans* isolates from different geographical areas is essential to design and produce an effective multivalent vaccine to prevent not only dental caries but also the prevention of life threatening diseases caused by *S. mutans*.
6. Lastly, our study also concludes that presence of cariogenic *S. mutans* is essential to initiate caries process in susceptible individuals with poor oral hygiene, sugar in diet and socioeconomically poor.

### Acknowledgement

Thanks to Gulbarga University, for the financial assistance during my tenure as PhD program, I thank

my Research guide, Prof.C.T.Shivannaveur. (Department of microbiology, research and studies) Gulbarga University. Gulbarga-585106.

### References

1. Balakrishnan M, Simmonds RS, Kilian M, Tagg JR. Different bacteriocin activities of *Streptococcus mutans* reflect distinct phylogenetic lineages. *J Med Microbiol* 2002; 51:941-948.
2. Hamada S, Ooshima T. Production and properties of bacteriocins (mutacins) from *Streptococcus mutans*. *Arch Oral Biol* 1975; 20:641-648.
3. Loesche WJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev* 2986; 50:353-380.
4. Clarke, J.K. On the bacterial factor in the etiology of dental caries. *Br J Exp Pathol* 1924; 5:141-147.
5. Banas, J.A. & Vickerman, M.M. Glucan-binding proteins of the oral streptococci. *Crit Rev Oral Biol Med* 2003; 14:89-99.
6. Bowen, W.H. & Koo, H. Biology of *Streptococcus mutans* derived glucosyltransferases: role in extracellular matrix formation of cariogenic biofilms. *Caries Res* 3022; 45:69-86.
7. Burne, R.A. Oral streptococci... products of their environment. *J Dent Res* 1998; 77:445-446.
8. Qi, F., Chen, P. & Caufield, P.W. Purification and biochemical characterization of mutacin I from the group I strain of *Streptococcus mutans*, CH43, and genetic analysis of mutacin I biosynthesis genes. *Appl Environ Microbiol* 2000; 66:3221-3229.
9. Qi et al., 2001; Kamiay et al., 2005): Regulation of *gbpC* expression in *Streptococcus mutans*. *Journal of Bacteriology*. 2007; 189: 6521-6531.

---

### Special Note!

Please note that our all Customers, Advertisers, Authors, Editorial Board Members and Editor-in-chief are advised to pay any type of charges against Article Processing, Editorial Board Membership Fees, Postage & Handling Charges of author copy, Purchase of Subscription, Single issue Purchase and Advertisement in any Journal directly to Red Flower Publication Pvt. Ltd. Nobody is authorized to collect the payment on behalf of Red Flower Publication Pvt. Ltd. and company is not responsible of respective services ordered for.