

Reduction of Cardiovascular Responses to Laryngoscopy and Intubation by Employing MgSO₄ Vs Normal Saline - A Comparative Study

C.K. Ramdas*, Deepak Falgunan**

Abstract

Background: Endotracheal intubation is often associated with a hypertension and tachycardia. Intravenous MgSO₄ is a popular method of blunting this response, because of its ability to depress sympathoadrenal response & catecholamine release.

Objective: it was undertaken to compare the effects of 50mg/kg MgSO₄ IV given 3 minutes before laryngoscopy and intubation.

Methods: A sample size of 60 patients aged 18 to 60 yrs were included in the study and they were allocated into 2 groups of n=30 each. Group I served as control. Group II received 50 mg/kg of IV magnesium sulphate 3 min before induction. HR, SBP were recorded pre-operatively, 30 sec, 1 min, 3min & 5min after intubation.

Results: Patient receiving iv MgSO₄ had a better intubating conditions (p<0.04) statistically significant than in Group I. There was an increase in the HR at 1 minute after intubation compared to basal value (p < 0.01). Also, a decrease in SBP observed at 1 minute and 3 min after intubation when compared with basal value (p<0.001) and (p<0.001) respectively. The decrease in RPP at 1 minute & 3 min after intubation when compared with the basal value (p < 0.001) & (p<0.01) respectively.

Conclusion: Magnesium sulphate 50 mg/kg IV infusion 3 minutes before induction, is a simple, effective and practical method of blunting cardiovascular responses to tracheal intubation, not associated with any adverse effect.

Keywords: Laryngoscopy; Tachycardia; Hypertension; Catecholamine.

Introduction

Direct laryngoscopy and endotracheal intubation following induction of anesthesia is almost always associated with hemodynamic changes due to reflex sympathetic discharge caused by epipharyngeal and laryngopharyngeal stimulation [1]. This increased sympathoadrenal activity may result in hypertension, tachycardia and arrhythmias [2-4]. This increase in blood pressure and heart rate are usually transitory, variable and unpredictable. Hypertensive patients are more prone to have significant increases in blood pressure whether they have been treated before hand or not. Transitory hypertension and tachycardia are probably of no consequence in healthy individuals but either or both may be hazardous to those with hypertension, myocardial insufficiency or cerebrovascular diseases. These

hemodynamic changes in such individuals may predispose to development of pulmonary oedema [5], myocardial insufficiency [6] and cerebrovascular accident [7]. At least in such individuals there is a necessity to blunt these harmful hemodynamic effects.

Many pharmacological methods have been devised to reduce the extent of hemodynamic events including high dose of opioids local anesthetics like lignocaine, alpha and beta adrenergic blockers and calcium channel blockers [1] vasodilatation drugs like nitro glycerine [8]. Topical anesthesia with lignocaine applied to the larynx and trachea in a variety of ways remains a popular method used alone or in combination with other techniques.

Intravenous MgSO₄ is a popular method of blunting this response, because of its ability to

Author's Affiliation:

*Assistant Professor, Department of Anesthesia, KMCT Medical College, Manassery-P.O, Mulkam, Kozhikode-673602, Kerala. **Assistant Professor, Department of Anesthesia, Kerala Medical College, Mangode, Cherpulassery, Palakkad Dt, Kerala -679503.

Corresponding Author:

C.K. Ramdas, Assistant Professor, Department of Anesthesia, KMCT Medical College, Manassery-P.O, Mulkam, Kozhikode-673602, Kerala state.

E-mail: thesisusm@gmail.com

Received on 12.12.2016

Accepted on 13.12.2016

depress sympathoadrenal response & catecholamine release.

Hence the present study was undertaken to compare the effect of Intravenous MgSO₄ and intravenous normal saline as a placebo on blunting the haemodynamic responses to endotracheal intubation.

Material and Methods

This study was a comparative clinical study of attenuation of cardiovascular responses to laryngoscopy and intubation, employing MgSO₄ vs Normal Saline was undertaken in MNR Medical College and Hospital, Sangareddy, during the June 2014 to July 2016. The study was undertaken after obtaining ethical committee clearance as well as informed consent from all the patients.

The study population were divided into two (2) sub groups with 30 patients in each group.

1. Control Group – Received normal saline as a placebo and served as control (n<30).
2. Study Group – Received 50 mg/kg of MgSO₄ i.v. 3 MIN before induction of anesthesia (n < 30).

Inclusion Criteria include patients belonging to ASA grades I and II and aged between 18 to 60 years undergoing various surgical procedure under general anaesthesia. Pre-anaesthetic evaluation and investigations was done on the evening before surgery. All patients were tested for any hypersensitivity reaction to local anaesthetics and an informed consent was obtained from all the patients. All the patients included in the study were premedicated with Tab. Alprazolam 0.5 mg and Tab. Ranitidine 150 mg orally at bed time the previous day.

Patients in control group received normal saline and served as control. Patients in study Group received Inj. MgSO₄ 50 mg/kg i.v. 3 minutes before induction of anesthesia. The intubating anesthetist was blinded to the study and dilutant procedure.

Anaesthesia was induced with inj. Thiopentone 5 mg/kg as 2.5% solution and endotracheal intubation was facilitated with succinylcholine 1.5 mg/kg administered one minute prior to laryngoscopy and intubation. The intubating conditions was evaluated and scoring was done according to the four step scale proposed by Goldberg and colleagues [9] by the intubating anaesthetist. It was graded as follow, Grade I- Excellent, Grade II- Good, Grade III- Poor,

Grade IV- Impossible intubating condition. The patients were intubated using appropriate sized cuffed endotracheal tubes. After confirming bilateral equal air entry, the endotracheal tube was secured.

Anaesthesia was maintained using 66% nitrous oxide and 33% of oxygen. After the patients recovered from succinylcholine further neuromuscular blockade was maintained with non-depolarizing muscle relaxants. At the end of the procedure patients were reversed with neostigmine 0.05 mg/kg IV and glycopyrrolate 0.008 mg/kg IV

The cardiovascular parameters were recorded in all the patients and cardiovascular parameters were noted as below

1. Pre-operative assessment before giving any study drugs and premedication
2. 30 seconds after intubation
3. 1 minute after intubation
4. 3 minute after intubation
5. 5 minute after intubation

The results were statistically evaluated using student 't' test & ANOVA test comparing between the groups and within the group respectively. P<0.05 was considered statistically significant.

Results

It is a prospective, controlled, randomized, double blind study to evaluate the efficacy of intravenous MgSO₄ 50 mg/kg and intravenous normal saline as a placebo on hemodynamic responses to laryngoscopy and endotracheal intubation.

Statistically there was no significant difference in two groups regarding the age (p>0.39), sex (p>1.00) and weight (p>0.33) (Table 1 and 2).

Regarding Changes in Heart Rate in Study Group:

The basal HR was 89.16±10.65 bpm, 1 minute after intubation, it was 97.20±18.82.. Subsequently, the elevated heart rate started settling down. By 3 minutes it was 96.60±18.11bpm and by 5 minutes it was 91.40±14.67 bpm. The increase in the HR at 1 minute after intubation compared to basal value was statistically highly significant (p < 0.01) (Table 3).

Changes in Systolic Blood Pressure in Study Group

The basal value of SBP was 127.86±10.27 mm Hg, 1 minute following intubation the SBP was

122.40±15.32 mm Hg. Afterwards the elevated blood pressure started coming down towards the baseline value.

By 3 minutes it was 119.50±10.90 mm Hg and by 5 minutes it was 117.53±9.34 mm Hg. The decrease in SBP observed at 1 minute and 3 min after intubation when compared with basal value was statistically significant (p < 0.001) and (p < 0.001) respectively

(Table 3).

Changes in Rate Pressure Product in Study Group

The basal RPP was 113.93±164.44. One minute after intubation, the RPP increased to 119.33±328.51. Subsequently the elevated RPP started settling down. By 3 minutes, it was 116.96±27.78 and by 5 minutes it was 107.20±185.24. The decrease in RPP 1 minute & 3 minute after intubation when compared with the

Table 1: Frequency distribution of cases according to age and weight

Parameter	Group	N	Mean ± SD	P value
AGE(years)	Group 1	30	35.0667±12.1199	0.39001
	Group 2	30	37.7333±11.7266	
Weight (kg)	Group 1	30	60.8667±7.9989	0.33939
	Group 2	30	63.1667±10.3460	

Table 2: Distribution of case according to gender

Group 1		Group 2	
Male	Female	Male	Female
18 (60%)	12 (40%)	18 (60%)	12(40%)

Table 3: Showing changes in mean heart rate, mean systolic bp and mean rate pressure product

Time	Group 1	Group 2
Mean Heart Rate		
Pre-operative	81.86±11.5	89.16±10.65
Intubation-30 sec	89.66±9.33	95.66±17.71
1 min	107.53±12.20	97.20±18.82
3 min	99.93±11.95	96.60±18.11
5 min	93.66±11.82	91.40±14.67
Mean Systolic Blood Pressure		
Pre-operative	122.66±11.72	127.86±10.27
Intubation- 30 sec	121.06±11.43	118.46±11.10
1 min	155.80±19.36	122.40±15.32
3 min	133.66±13.76	119.50±10.90
5 min	122.66±13.15	117.53±9.34
The Mean Rate Pressure Product (Rpp)		
Pre-operative	100.00±192.8	113.93±164.4
Intubation- 30 sec	108.26±161.30	113.30±271.6
1 min	168.26±326.52	119.33±321.5
3 min	134.33±254.43	11646.96±127.78
5 min	115.00±197.82	107.20±185.24

basal value was statistically significant (p < 0.001) & (p < 0.01) respectively (Table 3).

Discussion

Laryngoscopy and endotracheal intubation are associated with significant hypertension, tachycardia and arrhythmias. These hemodynamic responses were first recognised as early as in 1940 by Reid and Bruce et al [1]. They postulated that the disturbances in cardiovascular system were reflex in nature and mediated by the vagus nerve. In 1950 Burstein et al [2] studied the effects of laryngoscopy

and tracheal intubation on ECG changes and suggested the pressor response as consequences of an increase in sympathetic and sympathoadrenal activity.

These responses are transitory, variable and unpredictable and are much more pronounced in hypertensive patients than in normotensive individuals [4]. This hemodynamic stimulus is associated with increase in plasma nor-adrenaline concentrations parallel with the increase in blood pressure [3,9].

Michael F. M. et al [14] did a study to evaluate the effects of pre-treatment with 60 mg/kg body weight magnesium sulfate intravenous on cardiovascular

responses and catecholamine release associated with tracheal intubation were measured in 15 normal patients and in 15 saline solution pre-treated controls.

Magnesium pre-treatment increased heart rate by 13 ± 3.9 beats/minute. After intubation, heart rate was unchanged in the magnesium group at 107.3 ± 3.6 beats/minute but increased in the control group to 120.9 ± 4.6 beats/minute ($P < 0.05$). Systolic blood pressure increased after intubation from 106.8 ± 3.1 to 121.0 ± 4.4 mm Hg in patients given magnesium and from 106.4 ± 3.12 to 145.1 ± 5.6 mm Hg in the control group ($P < 0.05$).

It was concluded that magnesium sulfate attenuates the catecholamine mediated responses after tracheal intubation.

In our study, we found that SBP falls significantly after 1 minute ($p < 0.001$) & 3 minutes ($p < 0.001$) of intubation with decreased RPP after 1 minute ($p < 0.001$) & 3 minutes ($p < 0.01$) respectively in the study group. But at the same time, it is associated with considerable rise in the HR ($p < 0.01$) compared to the control group, thus showing similar results with the study.

Another study done by Yap LC [11] et al (1994) to evaluate the effect of magnesium sulfate on heart rate, blood pressure and hyperkalemic response following succinylcholine injection during tracheal intubation, using 60 mg/kg of magnesium sulfate intravenously 1min before succinylcholine injection. The control group received an equal volume of normal saline in the same way. Heart rate, blood pressure, venous and arterial potassium levels were measured at 1, 3, 5 and 10 minutes after intubation.

The results showed that magnesium sulfate could attenuate the hypertensive response at 1 minute and the hyperkalemic response at 1 and 3 minutes following succinylcholine-facilitated intubation; the tachycardia response at 1 minute after intubation could not be reduced by this agent.

The efficacy of intravenous magnesium to block the effects of sympathetic stimulation after nasotracheal intubation was studied by Jain PN et al (1995) [12]. They observed that IV magnesium sulphate in a dose of 60 mg/kg body wt. significantly ($p < 0.05$) attenuated the haemodynamic response to nasotracheal intubation without producing a significant rise in serum magnesium levels.

This study also got the same results of hemodynamic attenuation with iv magnesium sulfate, but associated with some amount of tachycardia after 1 min of intubation.

In our study, magnesium sulfate causes increase

in HR significantly after 1 min of intubation with significant reduction in SBP & RPP. After 3 min of intubation, further fall in SBP & RP occurred, thus showing consistency with the study [13]. However, we did not find any prolongation of neuromuscular blockade.

The hemodynamic variables were recorded in a study by G. D Puri et al [14] before induction, after the trial drug, after induction, and after endotracheal intubation. Magnesium sulfate administration was associated with increased cardiac index, a minimal increase in heart rate, and a significant decrease in mean arterial pressure (MAP) and systemic vascular resistance (SVR).

The magnesium group patients had a significantly lesser increase in MAP and SVR compared with control patients who received lidocaine before endotracheal intubation. Thus, magnesium is a useful adjuvant to attenuate endotracheal intubation response in patients with CAD.

However, in our study, we found that SBP falls significantly after 1 minute & 3 minutes of intubation with decreased RPP after 1 minute & 3 minutes respectively in the study group. But at the same time, it is associated with considerable rise in the HR compared to controls. Due to rise in HR, magnesium sulfate should be used with caution used for patients having coronary insufficiency or CAD.

In a randomized trial done by KH Naghibi et al [15] (2000) comparing the effects of magnesium sulphate with lidocaine for attenuating of presser response to tracheal intubations in 120 patients undergoing general anesthesia for cataract surgery. They have studied the effect of pretreatment with magnesium sulphate 50 mg/kg or 1.5 mg/kg lidocaine on this presser response found that there were no significant differences between two groups with respect preinduction of heart rate, mean arterial pressure, sex and age. A combined analysis showed that the magnesium sulphate were significantly better than 1.5mg/kg lidocaine in attenuating presser response to tracheal intubation, preventing postoperative tachycardia and hypertension and it is better than lidocaine.

Magnesium sulfate infusion in this study is effective in preventing increase in SBP & RPP after intubation but not effective in decreasing the heart rate. It was associated with tachycardia which come down to pre-operative level after 5 min of intubation.

Pipelzadeh MR et al [16] (2001) concluded that magnesium sulfate had a very limited usefulness in the attenuation of blood pressure and heart rate in

young healthy patients if given during induction of anesthesia. However, in this study, they had used magnesium sulfate in dose of 50 mg/ kg body wt just before laryngoscopy & found marked decrease in SBP & RPP after 1 & 3 minutes of intubation respectively.

Lim SH [17] et al (2007) studied the effects of magnesium sulfate and remifentanyl at attenuating the sympathetic responses were compared during laryngoscopy and endotracheal intubation.

Whether i.v. magnesium sulphate attenuates the haemodynamic stress responses to pneumoperitoneum by changing neurohumoral responses during laparoscopic cholecystectomy was studied by D. Jee et al [18] (2009), found that systolic and diastolic arterial pressures were greater in the control group (P<0.05) than in the magnesium group post-pneumoperitoneum, concluded that I.V. magnesium sulphate before pneumoperitoneum attenuates arterial pressure increases during laparoscopic cholecystectomy. This attenuation is apparently related to reductions in the release of catecholamine, vasopressin, or both.

In our study, we have found attenuation of SBP & RPP after intubation in magnesium sulfate group with slight increase in heart rate. This can be explained due to inherent property of drug in causing decreased catecholamine release [9] with vasodilatation thereby causing reduction in SBP & increase in heart rate.

Conclusion

We conclude that Magnesium sulphate 50 mg/kg IV infusion 3 minutes before induction, is a simple, effective and practical method of blunting cardiovascular responses to tracheal intubation, not associated with any adverse effect.

References

1. Reid LC and Bruce DE. Initiation of respiratory tract reflexes and its effects on heart. *Surg. Gynaec. Obstetrics*. 1940; 70:157.
2. Burstein CL, Lo Pinto FJ and Newman W. Electrocardiographic studies during endotracheal intubation 1, effects during usual routine techniques. *Anesthesiology*. 1950; 11:224.
3. King BD, Harris IC Jr., Freifenstein FE, Elder JD and Dripps RD. Reflex circulating responses to direct laryngoscopy and tracheal intubation performed during general anaesthesia. *Anesthesiology*. 1951; 12:556-566.
4. Prys-Roberts, Greene LT, Meloche R and Foex P. Studies of anaesthesia in relation to hypertension-II. Haemodynamic consequences of induction and endotracheal intubation. *British Journal of Anaesthesia*. 1971; 43:541-547.
5. Dalton B and Guiney T et al. Myocardial ischaemia from tachycardia and hypertension in coronary heart disease - Patient's undergoing anaesthesia. *Ann. Mtg. American Society of Anaesthesiologists, Boston*. 1972; 201-202.
6. Elisabeth J Fox, Garry S Sklar, Constance H Hill, Raymond Villanueva, Benton D King. Complication related to the pressor response to endotracheal intubation. *Anesthesiology*. 1977; 47:524-25.
7. Siedlecki J. Disturbances in the function of cardiovascular system in patients following endotracheal intubation and attempts of their prevention by pharmacological blockade of sympathetic system. *Anaesth Resusc Intensive Ther* 1975; 3:107-123.
8. van den Berg A A, Honjol D N M. Attenuation of the haemodynamic responses to noxious stimuli in patients undergoing cataract surgery. A comparison of magnesium sulphate, esmolol, lignocaine, nitroglycerine and placebo given i.v. with induction of anaesthesia. *European Journal of Anaesthesiology* 1997; 14:134-147.
9. Goldberg ME, Larijani GE, Azad SS, Sosis M, Seltzer JL, Ascher J et. Al comparison of tracheal intubating conditions and neuro muscular blocking profiles after intubating doses of mivvacurium chlorider or succinyl-choline in surgical outpatients. *Anesth Analg* 1989; 69:93-99.
10. Michael F. M., R. Eryk Beer, Jan D. Esser et al *Anesthesia & Analgesia*, 1989; 68(6):772-776.
11. Yap LC, Ho RT, Jawan B, Lee JH et al. Effects of magnesium sulfate pretreatment on succinylcholine-facilitated tracheal intubation *Acta Analgesiol Sin* 1994; 32(1):45-50.
12. Jain PN; Divatia JV, Manjure SS, Chattopadhyay G, Shah SC et al. Intravenous magnesium inhibits pressor response to nasotracheal intubation *Journal of Anaesthesiology Clinical Pharmacology*. 1995; 11(1):59-62.
13. Misuzu Nishiyama and Yuzuru Yanbe et al *Journal of anesthesia*, 1997; 11:18-21.
14. G D Puri, K S Marudhachalam, P Chari and R K Suri et al The effect of magnesium sulphate on hemodynamics and its efficacy in attenuating the response to endotracheal intubation in patients with coronary artery disease. *Anesthesia & Analgesia*, 1998; 87:808-811.
15. KH Naghibi, M Akhtari et attenuation of the pressor response to tracheal intubations by magnesium sulphate, *Journal of research in medical science*,

- 2000; 5(1).
16. Pipelzadeh MR, Poormehti Z, Mobasher M et al. Does Magnesium Sulfate attenuate hypertensive response to laryngoscopy and intubation? *Acta Medica Iranica* 2001; 39(3):150-152.
17. Lim SH, An DG, Choi SW, Lee SE, Kim YH, Lee JH, Lee KM, Cheong SH, Choi YK, Kim YJ, Shin CM The Comparison of Magnesium Sulfate and Remifentanyl
- in Attenuating Hemodynamic Response to Endotracheal Intubation et al. *Korean Journal of Anesthesiology* 2007; 53(5):577-582.
18. D. Jee, D. Lee, S. Yun and C. Lee et Magnesium sulphate attenuates arterial pressure increase during laparoscopic cholecystectomy al *Br. J. Anaesth.* 2009; 103(4):484-489.
-

Special Note!

Please note that our all Customers, Advertisers, Authors, Editorial Board Members and Editor-in-chief are advised to pay any type of charges against Article Processing, Editorial Board Membership Fees, Postage & Handling Charges of author copy, Purchase of Subscription, Single issue Purchase and Advertisement in any Journal directly to Red Flower Publication Pvt. Ltd. Nobody is authorized to collect the payment on behalf of Red Flower Publication Pvt. Ltd. and company is not responsible of respective services ordered for.